Hudsp to animals including, but not limited to, rabies . Animals appearing sick
appropriate authorities notified.

bones of the carcass. By dropping the bag over the carcass, and grabbing the carcass through the bag, any possibility of personal contamination can be completely avoided. A shovel or other tool may be used to lift and push the carcass into the bags.
    - Seal the bag with a knot or twist tie to avoid contaminating the vehicle. Avoid throwing animals into the truck bed to minimize splashing.
    - Deer and other large carcasses should be wrapped in plastic sheets or tarps
- The bag containing the carcass should be disposed of immediately in a campus trash dumpster.
- Gloves, tarps and any other materials should be bagged and disposed of
- The shovel, truck bed and any other contaminated non-disposable equipment should be disinfected using a 10% bleach solution. The shovel can be immersed in a bucket containing the solution and the solution can be sprayed or poured over the truck bed and any other surfaces. Contact the Custodial Supervisor to mix a fresh solution of 10% bleach. Nitrile/latex gloves and safety goggles should be worn while using the bleach solution.
- Hands should be washed well with soap and water after handling the carcass and after disinfecting equipment.

If you were bitten by an animal or had contact with the saliva or other fluids from an animal, **DO NOT** dispose of the carcass. Bag and save the carcass and refer to the Post Exposure Procedures, below.

TRAINING

All employees determined to have the potential for exposure will be provided appropriate training at the